


This article was downloaded by:[Meienberg, François]
On: 14 August 2007
Access Details: [subscription number 781322147]
Publisher: Informa Healthcare
Informa Ltd Registered in England and Wales Registered Number: 1072954
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Renal Failure

Publication details, including instructions for authors and subscription information:
<http://www.informaworld.com/smpp/title~content=t713597293>

A Fatal Case of Paraquat Poisoning Following Minimal Dermal Exposure

Online Publication Date: 01 March 2007

To cite this Article: Soloukides, Andreas, Moutzouris, Dimitrios-Anestis, Kassimatis, Theodore, Metaxatos, Georgios and Hadjiconstantinou, Valsamakis (2007) 'A Fatal Case of Paraquat Poisoning Following Minimal Dermal Exposure', Renal Failure, 29:3, 375 - 377

To link to this article: DOI: 10.1080/08860220601184134

URL: <http://dx.doi.org/10.1080/08860220601184134>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

© Taylor and Francis 2007

CASE REPORT

A Fatal Case of Paraquat Poisoning Following Minimal Dermal Exposure

Andreas Soloukides, Dimitrios-Anestis Moutzouris, Theodore Kassimatis, Georgios Metaxatos, and Valsamakis Hadjiconstantinou

Nephrology Department, “Evangelismos” General Hospital, Athens, Greece

Paraquat is a pesticide widely used in agriculture. Numerous cases of paraquat intoxication have been reported either accidentally or intentionally as a suicidal attempt. The most severe cases of paraquat poisoning refer to oral ingestion. Complications include respiratory, hepatic, and renal failure, and are usually fatal. Dermal exposure is less frequent and rarely fatal.

This article reports a case of an 81-year-old man with minimal skin burn after accidental paraquat exposure. The patient developed acute renal and respiratory failure and, despite aggressive treatment with hemodialysis, hemoperfusion, and mechanical ventilation, died two days later.

Keywords paraquat, pesticide, poisoning, renal failure, ARDS

INTRODUCTION

Paraquat (1,1'-dimethyl-4,4'-dipyridylum), the non-selective pesticide used worldwide in agriculture,^[1] is highly toxic in humans and has caused numerous fatalities. It remains the most common cause of fatal pesticide poisoning in western countries.^[2] Paraquat intoxication is characterized by circulatory failure with multiple organ failure in the early phase, as well as by respiratory failure caused by progressive pulmonary fibrosis in a later phase.^[3] Although severe cases of poisoning are usually the result of ingestion, accidental or in a suicide attempt, there are also sporadic reports of systemic involvement after skin exposure to paraquat.^[4,5]

This study reports a case of fatal paraquat poisoning after minimal dermal exposure.

Address correspondence to Dr. Andreas Soloukides, 23, Pontoeraklias Str, Athens 11527, Greece; Tel.: +30-210-7480448; Fax: +30-2107201624; E-mail: soloukides@yahoo.gr

CASE REPORT

An 81-year-old male presented to his family doctor because of a skin lesion of the right thigh after accidental contact with paraquat the previous evening. His relatives reported that the pesticide was spread on the trousers, and the old man slept overnight without removing the clothes. The lesion was limited, producing only skin erosion, which was treated empirically with steroid ointments. Four days later, the patient complained of severe breathlessness and was admitted in our department.

On admission, the patient complained of difficulty in breathing. He also reported decreased urine output over the last 48 hours. He did not report fever or other systemic symptoms. His past medical history was unremarkable except for mild hypertension during the last four years, which was treated with perindopril.

On initial physical examination, the patient was alert but highly distressed. Vital signs were (on admission): rectal temperature, 37.2°C; heart rate, 85/min; respiratory rate, 16/min; and blood pressure, 120/70 mmHg. Superficial chemical burn on the right thigh was calculated as covering 4% of the total body surface area. There were no mouth ulcers. The rest of the examination was negative.

Laboratory tests revealed the following: hematocrit, 42.5%; white blood cells count, $14.6 \times 10^3/\mu\text{L}$ (89.9% neutrophils); PLT, $160 \times 10^3/\mu\text{L}$; urea, 246 mg/dL; creatinine, 10.1 mg/dL; Na, 140 mEq/L; K, 4.2 mEq/L; alkaline phosphatase, 102 IU/L; SGOT, 87 IU/L; SGPT, 100 IU/L; LDH, 431 IU/L; Ca, 9.4 mg/dL; P, 6.0 mg/dL; ESR, 107 mm; PT, 26%; and INR, 2.17. Blood gas analysis on two liters of oxygen showed pH, 7.46; PaO₂, 49.1 mmHg; PaCO₂, 27.7 mmHg; and O₂ saturation 83.5%. Chest radiograph showed findings compatible with acute respiratory distress syndrome (ARDS; see Figure 1). Abdomen ultrasound was negative. Urine sodium dithionite test was negative for paraquat, and no pesticide was detected in blood sample. These findings indicated acute lung injury, acute renal injury, leucocytosis, and impaired hepatic function.



Figure 1. Chest x-ray showing infiltrates of the lower lobes, suggestive of Acute Respiratory Distress Syndrome (ARDS).

The patient was treated with hydration, oxygen supplementation, and intravenous antibiotics. Renal failure was managed by hemodialysis and hemoperfusion. The lung function continued to deteriorate. He was transferred to the Intensive Care Unit, intubated, and ventilated, but died two days later.

DISCUSSION

Paraquat poisoning remains a significant worldwide cause of morbidity and mortality,^[1] following pesticide exposure. The mechanism of toxicity in humans has not been clarified, but paraquat has been shown in animals to induce complete oxidation of both NADPH and NADH and increase lipid peroxidation.^[6] Additionally, paraquat is a potent generator of acute reactive oxygen species.^[6]

The most severe cases of paraquat poisoning occur following ingestion: oral exposure has been demonstrated to correlate with longer mean hospitalization and higher fatality rate.^[7] There are reports of dermal exposure to paraquat producing only local lesions on skin and nails without any signs of systemic involvement.^[8,9] Nevertheless, some cases of severe systemic involvement and deaths after skin exposure have been reported.^[4,5] In the present case, dermal exposure was extremely limited, covering only 4% of the total body surface area.

In vitro and in vivo experiments have indicated low percutaneous permeability of the substance through intact skin.^[10,11] Similar animal studies demonstrated that in an aqueous solution, paraquat ions do not penetrate the intact skin of the rabbit, which is more permeable than the human skin.^[12] Therefore, damaged skin is essential for

significant paraquat absorption.^[10,13] Previous skin lesions can be aggravated by paraquat itself. This was the case in the patient featured here, who remained in contact to paraquat through the humid trousers despite the initial chemical burn. Most fatalities via the dermal route are reported in developing countries.^[10,13] All cases are also associated with pre-existing skin lesions or prolonged contact of healthy skin with concentrated solutions.^[10,13]

There are many analyses in the literature that attempt to determine potent predictors of the outcome in paraquat poisoning. A multiple logistic regression analysis by Jones et al.^[14] demonstrated that patients with a given concentration of paraquat have a better chance of survival if they have only recently taken it; moreover, at any time after ingestion, the prognosis is better the lower the plasma concentration. As kidneys are the major organ to excrete paraquat, paraquat-poisoned patients with severe acute renal failure have a prolonged half-life of paraquat and a markedly increased mortality rate.^[15]

Retrospective studies showed that good predictors of outcome may be the low plasma and urine concentrations within the first 24 hours of intoxication.^[16,17] Nevertheless, the qualitative urine test is not reliable for patients presenting >24 hours after the exposure.^[17]

Treatment choices for paraquat poisoning include general treatment of poisoning (aiming to decontaminate the gut and minimize absorption), hypo-oxygenation,^[3] lung radiotherapy, hemodialysis, and hemoperfusion,^[3] but results are controversial.^[18,19] Nevertheless, we decided to treat our patient with hemoperfusion, as there are anecdotal reports of successful treatment even in cases of late referral.^[20] There are also reports with promising results regarding the use of combined methylprednisolone and dexamethazone therapy.^[21,22] There is one report in the literature for the successful treatment of severe paraquat poisoning after dermal exposure by the combined use of cyclophosphamide and methylprednisolone pulse therapy.^[23]

Steroid treatment may decrease the inflammation of paraquat poisoning.^[24] Moreover, steroids can suppress superoxide production by neutrophils.^[25] Therefore, steroid therapy may be an effective treatment of paraquat poisoning (after the failure of standard therapies), where the inflammation of the lungs may play a major role in the lethal hypoxemia of patients.^[22]

In conclusion, it is clear that even minimal dermal exposure to paraquat can be fatal, especially when associated with high concentrations of the substance and/or significant delay in treatment initiation. The exact efficacy of the current medical management remains controversial. Therefore, it is crucial for emergency physicians to suspect paraquat poisoning when facing chemical burns and skin lesions and to investigate for the possibility of intoxication, particularly when skin lesions are associated with systemic symptoms.

REFERENCES

- Jeyaratnam J. Acute pesticide poisoning: a major global health problem. *World Health Stat Q*. 1990;43(3):139–144.
- Thompson JP, Casey PB, Vale JA. Deaths from pesticide poisoning in England and Wales 1990–1991. *Hum Exp Toxicol*. 1995;14(5):437–445.
- Vale JA, Meredith TJ, Buckley BM. Paraquat poisoning: clinical features and immediate general management. *Hum Toxicol*. 1987;6(1):41–47.
- Papiris SA, Maniati MA, Kyriakidis V, Constantopoulos SH. Pulmonary damage due to paraquat poisoning through skin absorption. *Respiration*. 1995;62(2):101–103.
- Tungsanga K, Chusilp S, Israsena S, Sitprija V. Paraquat poisoning: evidence of systemic toxicity after dermal exposure. *Postgrad Med J*. 1983;59(691):338–339.
- Togashi H, Shinzawa H, Wakabayashi H, et al. Superoxide is involved in the pathogenesis of paraquat-induced injury in cultured rat liver slices. *Hepatology*. 1991;14(4 Pt 1):707–714.
- Lin CL, Yang CT, Pan KY, Huang CC. Most common intoxication in nephrology ward organophosphate poisoning. *Ren Fail*. 2004;26(4):349–354.
- Bismuth C, Garnier R, Dally S, Fournier PE, Scherrmann JM. Prognosis and treatment of paraquat poisoning: a review of 28 cases. *J Clin Toxicol*. 1982;19(5):461–474.
- Fitzgerald GR, Barniville G, Black J, Silke B, Carmody M, O'Dwyer WF. Paraquat poisoning in agricultural workers. *Ir Med J*. 1978;71(10):336–342.
- Garnier R. *Paraquat poisoning by inhalation or skin absorption*. New York: Marcel Dekker; 1995.
- Wester RC, Mailbach HI, Bucks DA. Paraquat poisoning by skin absorption. *Hum Toxicol*. 1989;8(3):251–252.
- Guardascione V. A contribution to the knowledge of professional pathology from paraquat, a herbicide based on dipyridyl. *Folia Med (Napoli)*. 1969;52:728–738.
- Smith JG. Paraquat poisoning by skin absorption: a review. *Hum Toxicol*. 1988;7(1):15–19.
- Jones AL, Elton R, Flanagan R. Multiple logistic regression analysis of plasma paraquat concentrations as a predictor of outcome in 375 cases of paraquat poisoning. *QJM*. 1999;92(10):573–578.
- Dabir D, Russell LN, Ronald DF. Nephrotoxicity of paraquat in man. *Arch Intern Med*. 1979;139:172–174.
- Braithwaite RA. Emergency analysis of paraquat in biological fluids. *Hum Toxicol*. 1987;6(1):83–86.
- Scherrmann JM, Houze P, Bismuth C, Bourdon R. Prognostic value of plasma and urine paraquat concentration. *Hum Toxicol*. 1987;6(1):91–93.
- Hampson EC PS. Failure of haemoperfusion and haemodialysis to prevent death in paraquat poisoning. A retrospective review of 42 patients. *Med Toxicol Adverse Drug Exp*. 1988;3(1):64–71.
- Bohler JWR, Keller E, Logemann E, Just H, Schollmeyer PJ. Continuous arteriovenous haemoperfusion (CAVHP) for treatment of paraquat poisoning. *Nephrol Dial Transplant*. 1992;7:875–878.
- Katopodis K, Logothetis E, Noussias C, Hadjiconstantinou V. Survival of a paraquat poisoned patient, despite late (four days) referral and initiation of conventional haemoperfusion treatment. *Nephrol Dial Transplant*. 1993;8:570–571.
- Perriens JH, Benimadho S, Kiauw IL, Wisse J, Chee H. High-dose cyclophosphamide and dexamethasone in paraquat poisoning: a prospective study. *Hum Exp Toxicol*. 1992;11(2):129–134.
- Chen GH, Lin JL, Huang YK. Combined methylprednisolone and dexamethasone therapy for paraquat poisoning. *Crit Care Med*. 2002;30(11):2584–2587.
- Lin NC, Lin JL, Lin-Tan DT, Yu CC. Combined initial cyclophosphamide with repeated methylprednisolone pulse therapy for severe paraquat poisoning from dermal exposure. *J Clin Toxicol*. 2003;41(6):877–881.
- Lin JL, Leu ML, Liu YC, Chen GH. A prospective clinical trial of pulse therapy with glucocorticoid and cyclophosphamide in moderate to severe paraquat-poisoned patients. *Am J Respir Crit Care Med*. 1999;159(2):357–360.
- Youshida TTM, Sotomatsu A, et al. Effect of methylprednisolone pulse therapy on superoxide production of neutrophils. *Neurol Res*. 1999;21:509–512.

