



UNEP

**United Nations  
Environment Programme**



**Food and Agriculture Organization  
of the United Nations**

Distr.: General  
1 December 2005

English only

**Rotterdam Convention on the Prior Informed  
Consent Procedure for Certain Hazardous  
Chemicals and Pesticides in International Trade  
Chemical Review Committee**

Second meeting

Geneva, 13–17 February 2006

Item 5 (b) of the provisional agenda\*

**Inclusion of chemicals in Annex III of the Rotterdam Convention:  
review of notifications of final regulatory actions to ban  
or severely restrict a chemical: Endosulfan**

## **Endosulfan: additional supporting documentation provided by Thailand**

### **Note by the secretariat**

The annex to the present note contains additional supporting documentation provided by Thailand in support of its final regulatory action on endosulfan.

---

\* UNEP/FAO/RC/CRC.2/1.

## Annex

### List of additional supporting documentation on Endosulfan submitted by Thailand

- Effects of Endosulfan on Male Reproductive Development (Saiyed H., Dewan.A et al). Environmental Health Perspectives Volume 111 Number 16, December 2003
  - Slowing Puberty? Pesticides may hinder development in boys. Kate Ramsayar. Science News Online (week of December 13, 1003, Vol. 164, No. 24)
  - Endosulfan: Public Health Fact Sheets (Queensland Government).
-

## Children's Health

### Effect of Endosulfan on Male Reproductive Development

Habibullah Saiyed,<sup>1</sup> Aruna Dewan,<sup>1</sup> Vijay Bhatnagar,<sup>1</sup> Udyavar Shenoy,<sup>2</sup> Rathika Shenoy,<sup>2</sup> Hirehall Rajmohan,<sup>3</sup> Kumud Patel,<sup>1</sup> Rekha Kashyap,<sup>1</sup> Pradip Kulkarni,<sup>1</sup> Bagalur Rajan,<sup>3</sup> and Bhadabhai Lakkad<sup>1</sup>

<sup>1</sup>National Institute of Occupational Health (Indian Council of Medical Research), Meghani Nagar, Ahmedabad, India; <sup>2</sup>Department of Pediatrics, Kasturba Medical College, Mangalore, India; <sup>3</sup>Regional Occupational Health Research Centre, Bangalore, India

#### Abstract

There is experimental evidence of adverse effects of endosulfan on the male reproductive system, but there are no human data. Therefore, we undertook a study to examine the relationship between environmental endosulfan exposure and reproductive development in male children and adolescents. The study population was composed of 117 male schoolchildren (10-19 years of age) of a village situated at the foothills of cashew plantations, where endosulfan had been aerially sprayed for more than 20 years, and 90 comparable controls with no such exposure history. The study parameters included recording of clinical history, physical examination, sexual maturity rating (SMR) according to Tanner stages, and estimation of serum levels of testosterone, luteinizing hormone (LH), follicle-stimulating hormone, and endosulfan residues (70 study and 47 control subjects). Mean  $\pm$  SE serum endosulfan levels in the study group ( $7.47 \pm 1.19$  ppb) were significantly higher ( $p < 0.001$ ) than in controls ( $1.37 \pm 0.40$  ppb). Multiple regression analysis showed that SMR scoring for development of pubic hair, testes, penis, and serum testosterone level was positively related to age and negatively related to aerial exposure to endosulfan (AEE;  $p < 0.01$ ). Serum LH levels were significantly positively related to AEE after controlling for age ( $p < 0.01$ ). The prevalence of congenital abnormalities related to testicular descent (congenital hydrocele, undescended testis, and congenital inguinal hernia) among study and controls subjects was 5.1% and 1.1%, respectively, but the differences were statistically nonsignificant. Our study results suggest that endosulfan exposure in male children may delay sexual maturity and interfere with sex hormone synthesis. Our study is limited by small sample size and nonparticipation. *Key words:* endocrine disruptor, endosulfan, luteinizing hormone, male reproductive development, sexual maturity rating, testosterone. *Environ Health Perspect* 111:1958-1962 (2003). doi:10.1289/ehp.6271 available via <http://dx.doi.org/> [Online 22 September 2003]

---

Address correspondence to H.N. Saiyed, Director, National Institute of Occupational Health, Meghani Nagar, Ahmedabad 380016, India. Telephone: 91-79-268-6351. Fax: 91-79-268-6110. E-mail: [saiyedhn@yahoo.com](mailto:saiyedhn@yahoo.com)

We gratefully acknowledge the Ministry of Health and Family Welfare, Government of India, for sponsoring this study; volunteers who participated in the study; and the pediatricians from Department of Pediatrics, Kasturba Medical College, Mangalore, India.

The authors declare they have no competing financial interests.

Received 10 February 2003; accepted 22 September 2003.

The full version of this article is available for free in [HTML](#) or [PDF](#) formats.

*Environmental Health Perspectives* Volume 111, Number 16, December 2003

## Children's Health

# Effect of Endosulfan on Male Reproductive Development

Habibullah Saiyed,<sup>1</sup> Aruna Dewan,<sup>1</sup> Vijay Bhatnagar,<sup>1</sup> Udyavar Shenoy,<sup>2</sup> Rathika Shenoy,<sup>2</sup> Hirehall Rajmohan,<sup>3</sup> Kumud Patel,<sup>1</sup> Rekha Kashyap,<sup>1</sup> Pradip Kulkarni,<sup>1</sup> Bagalur Rajan,<sup>3</sup> and Bhadabhai Lakkad<sup>1</sup>

<sup>1</sup>National Institute of Occupational Health (Indian Council of Medical Research), Meghani Nagar, Ahmedabad, India;

<sup>2</sup>Department of Pediatrics, Kasturba Medical College, Mangalore, India; <sup>3</sup>Regional Occupational Health Research Centre, Bangalore, India

- [Introduction](#)
- [Materials and Methods](#)
- [Results](#)
- [Discussion](#)

### Abstract

There is experimental evidence of adverse effects of endosulfan on the male reproductive system, but there are no human data. Therefore, we undertook a study to examine the relationship between environmental endosulfan exposure and reproductive development in male children and adolescents. The study population was composed of 117 male schoolchildren (10-19 years of age) of a village situated at the foothills of cashew plantations, where endosulfan had been aerially sprayed for more than 20 years, and 90 comparable controls with no such exposure history. The study parameters included recording of clinical history, physical examination, sexual maturity rating (SMR) according to Tanner stages, and estimation of serum levels of testosterone, luteinizing hormone (LH), follicle-stimulating hormone, and endosulfan residues (70 study and 47 control subjects). Mean  $\pm$  SE serum endosulfan levels in the study group ( $7.47 \pm 1.19$  ppb) were significantly higher ( $p < 0.001$ ) than in controls ( $1.37 \pm 0.40$  ppb). Multiple regression analysis showed that SMR scoring for development of pubic hair, testes, penis, and serum testosterone level was positively related to age and negatively related to aerial exposure to endosulfan (AEE;  $p < 0.01$ ). Serum LH levels were significantly positively related to AEE after controlling for age ( $p < 0.01$ ). The prevalence of congenital abnormalities related to testicular descent (congenital hydrocele, undescended testis, and congenital inguinal hernia) among study and controls subjects was 5.1% and 1.1%, respectively, but the differences were statistically nonsignificant. Our study results suggest that endosulfan exposure in male children may delay sexual maturity and interfere with sex hormone synthesis. Our study is limited by small sample size and nonparticipation. **Key words:** endocrine disruptor, endosulfan, luteinizing hormone, male reproductive development, sexual maturity rating, testosterone. *Environ Health Perspect* 111:1958-1962 (2003). doi:10.1289/ehp.6271 available via <http://dx.doi.org/> [Online 22 September 2003]

---

Address correspondence to H.N. Saiyed, Director, National Institute of Occupational Health, Meghani Nagar, Ahmedabad 380016, India. Telephone: 91-79-268-6351. Fax: 91-79-268-6110. E-mail: [saiyedhn@yahoo.com](mailto:saiyedhn@yahoo.com)

We gratefully acknowledge the Ministry of Health and Family Welfare, Government of India, for sponsoring this study; volunteers who participated in the study; and the pediatricians from Department of Pediatrics, Kasturba Medical College, Mangalore, India.

The authors declare they have no competing financial interests.

Received 10 February 2003; accepted 22 September 2003.

## Introduction

Endosulfan (6,7,8,9,10,10-hexachloro-1,5,5a,6,9,9a-hexahydro-6,9-methano-2,4,3-benzodioxathiepin-3-oxide) is a broad-spectrum insecticide and acaricide first registered for use in the United States in 1954 to control agricultural insect and mite pests on a variety of field, fruit, and vegetable crops. Technical-grade endosulfan is composed of two stereochemical isomers,  $\alpha$ -endosulfan and  $\beta$ -endosulfan, in concentrations of approximately 70% and 30%, respectively. Use data from 1987 to 1997 indicate an average domestic use of approximately 1.38 million pounds of active ingredient per year [U.S. Environmental Protection Agency (U.S. EPA) 2002]. It has been found in at least 162 of the 1,569 current National Priorities List sites by the U.S. EPA (HazDat 2000). In India, it is widely used against a variety of agricultural pests. During 1999-2000, about 81,000 metric tons of endosulfan was manufactured in India, and in terms of tonnage its production was next only to mancozeb (103,000 metric tons) and monocrotophos (95,000 metric tons) (Anonymous 2001).

Oral LD50 (lethal dose sufficient to kill 50% of population) endosulfan in rats is 80 mg/kg, and it has been classified as a moderately hazardous (class II) pesticide [World Health Organization (WHO) 2002]. Neurotoxicity is the major end point of concern in acute endosulfan exposure in human beings and experimental animals. No data are available for subacute or chronic exposure to endosulfan in human subjects; however, the subacute and chronic toxicity studies of endosulfan in animals suggest that the liver, kidneys, immune system, and testes are the main target organs [Agency for Toxic Substances and Disease Registry (ATSDR) 2000].

In recent years, there has been growing concern about toxicity of a number of chemicals, including pesticides, on the male reproductive system (Murray et al. 2001; Sharpe 2001). Reported effects of endosulfan on the male reproductive system in experimental animals have been variable, depending on species, age at exposure, dose, duration of exposure, and study end points. Routine gross and histopathologic examination of the reproductive organs of male mice that consumed doses of 7.3 mg/kg/day for 13 weeks (Hoechst. Unpublished data) or 2.5-5.0 mg/kg/day for 2 years [Hack et al. 1995; Hoechst. Unpublished data; National Cancer Institute (NCI) 1978] revealed no toxic effects. Later on, more detailed studies in adult rats exposed to 2.5, 5, and 10 mg/kg/day endosulfan for 5 days per week for 10 weeks showed reduced intratesticular spermatid counts, sperm abnormalities, and changes in the marker enzymes of testicular activities, such as lactate dehydrogenase, sorbitol dehydrogenase,  $\gamma$ -glutamyl transpeptidase, and glucose-6-phosphate dehydrogenase, providing further evidence of effects on spermatogenesis (Khan and Sinha 1996; Sinha et al. 1995). Exposure of younger animals (3 weeks old) showed marked depletion of spermatid count as well as decreased daily sperm production at a dose of 2.5 mg/kg/day (Sinha et al. 1997), which was earlier seen only at 5 mg/kg/day in adult rats by the same investigators (Sinha et al. 1995). More recent studies have shown that exposure of pregnant rats to endosulfan at 1 mg/kg/day from day 12 through parturition leads to decreased spermatogenesis in offspring (Sinha et al. 2001). Dalsenter et al. (1999) reported similar observations at 3 mg/kg/day but not at 1.5 mg/kg/day, and they attributed this to strain variation (Dalsenter et al. 2003). Thus, experimental studies suggest that endosulfan can affect the male reproductive system and also that these effects are likely to be greater if exposure occurs during the developmental phase.

Environmental exposure to a single chemical over a long period of time is very rare. We came across a situation where endosulfan was the only pesticide that had been aerially sprayed two to three times a year for more than 20 years on cashew nut plantations situated on hilltops in some villages of northern Kerala, India (Figure 1). The population living in the valley had a significant chance of exposure to this pesticide during aerial spray and subsequently through other contaminated environmental media. This population, therefore, provided a unique opportunity to study the long-term health effects of endosulfan. In this article, we report the effects of endosulfan on male reproductive development.

## Materials and Methods

**Selection of study and control areas.** The exposed population was defined as schoolchildren who were permanent residents of the village situated below the cashew plantations where endosulfan had been sprayed aerially. This village had 12 first-order streams originating from the cashew plantations. Most of the habitations were along the valleys and close to the stream banks. Most of the inhabitants depended on runoff water for irrigation purposes. The control population was selected from schoolchildren of another village situated approximately 20 km away. The population of this village was comparable with the exposed population in socioeconomic status, ethnicity, and occupational characteristics but without any history of aerial endosulfan spray. The control village did not have streams.

**Selection of study and control subjects.** The main study was carried out to explore the effects of endosulfan exposure on growth and development in 619 schoolchildren of both sexes (5-19 years) and 416 comparable controls. All male children (272 exposed and 135 controls) older than 10 years were asked to participate in a sexual maturity rating (SMR) study; 117 (43%) exposed and 90 (67%) controls participated in SMR examination. For the hormone study, every other student who participated in the SMR study was requested to donate a blood sample.

**Study parameters.** The study parameters included recording of clinical history in a specially designed pro forma physical examination, assessment of SMR by Tanner's classification (Marshall and Tanner 1969), and estimation of serum levels of testosterone, luteinizing hormone (LH), follicle-stimulating hormone (FSH), and endosulfan residues.




Figure 1. Cashew nut plantations on the hills and houses in the valley. Water streams formed in the hills pass through residential zones in the valley.




Figure 2. Chromatograms of  $\alpha$ -endosulfan,  $\beta$ -endosulfan, and endosulfan sulfate. RT, retention time. (A) Standard chromatogram of  $\alpha$ -endosulfan,  $\beta$ -endosulfan, and endosulfan sulfate. (B)

**Ethical aspects.** This study was approved by the ethics committee of the National Institute of Occupational Health. Parents, who were requested to accompany their children at the time of examination, were told the objectives of the study, and a consent form in local language was read aloud to them. The children were examined only after one of the parents gave written consent. In addition, special consent of the child was taken for the SMR study, and only in willing cases were blood samples collected. The SMR examination was carried out by pediatricians observing necessary privacy required for this delicate examination.

**Collection, storage, and transport of blood samples.** Five milliliters of venous blood were collected from each willing individual between 1000 and 1200 hr on the day of examination and was centrifuged at 5,000 rpm for 5 min in the field laboratory. Serum was separated and stored at -20°C in a nearby hospital. The serum samples were air-shipped under dry ice to the laboratories at National Institute of Occupational Health, Ahmedabad, India. To avoid observer bias, the samples were coded before being handed over for analysis.

**Chemicals and standard control materials for analysis of endosulfan in serum samples.** All the chemicals and reagents used in the extraction and cleanup of endosulfan residues were highly pure HPLC (high-performance solvents filtered through 0.2-µ filters and packed under nitrogen)-grade obtained from Qualigens Fine Chemicals (Glaxo India Ltd., Mumbai, India) and were checked for any pesticide contamination. Glassware used was free from residue contamination. Standard reference materials of α-endosulfan (99.0%), β-endosulfan (99.0%), and endosulfan sulfate (99.0%) were a gift from M/s Excel Ltd., (Mumbai, India), which is the largest manufacturer of endosulfan in India.

**Extraction of residues from serum.** Extraction was modified from techniques described in Dale et al. (1966) and U.S. EPA (1980). Briefly, serum (0.5 mL) was pipetted into a graduated stoppered centrifuge tube, 6 mL of hexane was added, and it was rotated in a slow-speed Roto-rac (National Institute of Occupational Health, Ahmedabad, India) for 2 hr. The organic layer was transferred into another graduated tube and was evaporated to dryness under a stream of nitrogen. The final volume was made up with hexane corresponding to the expected concentration of the residue. A suitable aliquot was injected into a gas chromatograph with an electron capture detector. We calculated the recoveries of endosulfan residues, which ranged from 88 to 102%. In addition, fortified samples were studied as a part of quality assurance and quality control.

**Instrumentation and quantification.** We used a gas chromatograph (model 6890) equipped with a Micro Electron Capture Detector, a capillary column (HP 5, 60 m, 0.25 mm inner diameter, film thickness 0.25 µm; all these items from Hewlett Packard Agilent Plus; Agilent Technologies, Little Fall, DE, USA), and N<sub>2</sub> (ultra high purity, 99.999% grade) as carrier gas for the quantification of endosulfan residues. The initial oven temperature was 80°C with ramp rate of 20°C per min to 200°C. The injector port temperature was 220°C (splitless mode), and detector temperature was 275°C. We quantified the samples by comparing the peak area of each with those of their respective standards. The retention times of α-endosulfan, β-endosulfan, and endosulfan sulfate were at 38.9, 52.5, and 67.2 min, respectively (Figure 2A,B).

**Hormone analyses.** We estimated testosterone, LH, and FSH in 50 µL and 100 µL serum samples of study and control subjects using radioimmunoassay kits procured from Immunotech (Marseille, France). We used a Wallac 1470 Wizard autogamma counter (Perkin Elmer, Turku, Finland) to count radioactivity with detection efficiency of 78% for <sup>125</sup>I and negligible cross talk with other isotopes. Equipment and glassware were segregated to prevent cross-contamination. The hormones were estimated in serial dilutions of serum along with parallel curves to standard. All the samples for hormone estimations were

Chromatogram of α-endosulfan, β-endosulfan, and endosulfan sulfate in blood sample of a study subject.

Table 1.

Table 1. Comparison of the mean (± SD) age and some of the growth-related basic parameters in study and control subjects participating in the SMR examination.

Parameters	Control (n=90)	Study (n=117)
Age (years)	13.10 ± 2.12	12.80 ± 2.07
Height (cm)	141 ± 10.60	139 ± 13.30
Weight (kg)	30.70 ± 7.44	29.50 ± 8.93
Body mass index	15.20 ± 1.98	15.00 ± 2.11
Skin-fold thickness (mm)	7.31 ± 2.15	7.40 ± 2.28

Table 2.

Table 2. Summary of the multiple regression analyses of endosulfan (µg/ml) against age and AEE. Regression models are adjusted for AEE and age.

Endosulfan isotope	R <sup>2</sup>	F-value	p-value	Age	AEE
α-Endosulfan	0.16**	2.64**	0.11**	0.03	-0.03**
β-Endosulfan	0.15**	2.58**	0.12**	0.05	-0.02*
Endosulfan sulfate	0.12**	2.28**	0.13**	0.02	-0.02**

No. of subjects: α-Endosulfan = 117; β-Endosulfan = 117; Endosulfan sulfate = 117.




Figure 3. SMR score for the development of pubic hair according to age and AEE.




Figure 4. SMR score for the development of testes according to age and AEE.



processed in one assay to rule out interassay variations. We performed a linearity study to assess the sensitivity of the hormone assays by serial dilution of a high level with the zero standards. The sensitivity to testosterone, LH, and FSH was 2.5 ng/dL, 0.45 IU/L and 0.02 IU/L, respectively.

**Statistical analysis.** The statistical software SPSS, release 6.1.4 (SPSS Inc., Chicago, IL, USA), was used for statistical analysis of data and for making graphs. For the multiple regression analysis, age and aerial exposure to endosulfan (AEE; study = 1, control = 0) were taken as independent variables, and parameters of SMR and serum hormone levels were taken as dependent variables. Exact age in years on the date of examination was used as the age variable.

We used the following multiple regression equation: SMR score =  $b_0 + b_1 \text{ age} + b_2 \times \text{AEE}$ , where  $b_0$  is the regression constant and  $b_1$  and  $b_2$  are the regression coefficients of age and exposure, respectively, fitted for SMR scores of pubic hair, testes, and penis. Similarly, multiple regression equations were fitted for testosterone, LH, and FSH levels.

**Results**

Table 1 shows the comparison of the mean  $\pm$  SD values of the age and some of the growth-related basic parameters in study and control subjects participating in the SMR study. There are no statistically significant differences in the mean values of these parameters in two groups. The mean values of height, weight, body mass index, and skin-fold thickness of the nonparticipating subjects in study and control groups with respect to age were also comparable, indicating that the nonparticipation of the subjects may not have had significant effects on the outcome of the study.

Clinical examination revealed six (5.1%) cases of congenital malformations in the study group, consisting of undescended testis (two), congenital hydrocele (three), and congenital inguinal hernia (one), and only one case (1.1%) of congenital inguinal hernia in the control group. The differences in prevalence of congenital malformation were statistically nonsignificant.

**SMR score.** A summary of the multiple regression analysis of parameters of SMR score on age and AEE is presented in Table 2. The values of  $R^2$  corresponding to SMR of pubic hair, testes, and penis are 0.48, 0.43, and 0.43, respectively, are statistically significant ( $p < 0.001$ ), and indicate that considerable proportion of variance of SMR scores can be attributed to age and AEE. The statistically significant negative regression coefficient ( $b_2$ ) of AEE in all three equations signifies delayed sexual maturity in the study group compared with the control group. As expected, age had a significant positive regression coefficient.


These effects of AEE are further elaborated graphically in Figures 3-5. In Figure 3, the findings for SMR for pubic hair is plotted against age for study and control individuals. The regression lines drawn for the study and control groups indicate that the SMR score for the study group is lower for the same age. Similar observations were made for the SMR score for testicular (Figure 4) and penis development (Figure 5).

**Serum hormone levels.** Table 3 shows the summary of the multiple regression analysis of serum testosterone levels against age, AEE, and serum LH levels. The value of  $R^2$  is 0.61, which means that 61% of the variations observed in serum testosterone levels can be explained on the basis of age, AEE, and serum LH levels. Very small  $p$ -values ( $< 0.001$  in all cases) of the regression coefficient indicate that statistically these parameters are significant in determining the serum testosterone levels. The positive sign of the regression coefficient for age and LH indicates that with increase in age and serum LH levels, there is an increase in the serum testosterone levels. This is a well-known physiologic fact. On the other hand, the regression coefficient for AEE has a negative sign, which means the serum testosterone levels of individuals belonging to the study group are statistically lower than expected from age and serum LH levels. The levels of FSH were slightly higher for the age in study group; however, the differences were statistically nonsignificant.


**Figure 5.** SMR score for the development of the penis according to age and AEE.

**Table 3.**

Parameter	Study (n=45)	Control (n=70)
Intercept	0.11**	0.05
Age (years)	0.01**	0.01**
AEE	-0.001**	0.0002**
Serum LH (IU/L)	0.0001**	0.0001**



**Figure 6.** Serum testosterone levels according to age and AEE.



**Figure 7.** Serum LH levels according to age and AEE.

**Table 4.**

	Control (n=45)	Study (n=70)
$\alpha$ -Endosulfan	0.07 $\pm$ 0.23	4.24 $\pm$ 0.74**
$\beta$ -Endosulfan	0.10 $\pm$ 0.17	1.77 $\pm$ 0.38**
Endosulfan sulfate	0.10 $\pm$ 0.08	1.47 $\pm$ 0.33**
Total endosulfan	1.37 $\pm$ 0.40	7.47 $\pm$ 1.19**

\*\*  $p < 0.001$ .

In Figure 6, serum testosterone levels are plotted against age for study and control individuals. Regression lines drawn for the study and control individuals indicate that the average serum testosterone levels for the study group are lower for the same age. Similarly, in Figure 7, serum LH levels are plotted against age in study and control individuals. The regression lines indicate that average serum levels for the same age are higher in the study group.

**Endosulfan exposure.** Endosulfan was detected in serum samples of 78% of the children in the study group and 29% of the children in the control group. Table 4 shows the serum endosulfan levels in the study and control groups. The levels of endosulfan in the study group children are significantly higher ( $p < 0.001$ ).

## Discussion

Our study results, after controlling for age, showed significantly lower SMR scores and serum testosterone levels and higher levels of serum LH in the study group compared with controls. To link these changes with endosulfan exposure, we should look at two issues: biologic plausibility of the cause-effect relationship and pathways of endosulfan exposure.

**Biologic plausibility.** There are reports of testicular toxicity of endosulfan manifested as decreased spermatogenesis and testicular hormone synthesis (steroidogenesis), as evidenced by a decrease in spermatid count in testes and in sperm count in the cauda epididymis and by changes in marker enzymes for testicular steroidogenesis in adult animals (Chitra et al. 1999; Singh and Pandey 1989, 1990; Sinha et al. 1995). These effects were seen at much lower dosages and shorter durations if exposures occurred during the prenatal or prepubertal periods (Dalsenter et al. 1999; Sinha et al. 1995, 1997, 2001). Singh and Pandey (1990) also reported profound decreases in the levels of plasma LH, FSH, and testosterone associated with decrease in testicular testosterone in pubertal rats exposed to endosulfan for 30 days. Thus, our observations of low testosterone levels in male children conform with the animal studies. Lower SMR scores appear to reflect lower serum testosterone levels for age. In our study, it is not possible to confirm disturbed spermatogenesis observed in animal studies.

The higher prevalence of congenital abnormalities related to testicular descent observed in the study group should not be overlooked simply because it failed to achieve statistical significance (which may be due to small sample size), because there is indirect evidence of endosulfan exposure associated with undescended testes in a human population from Spain. Garcia-Rodriguez et al. (1996) reported a higher incidence of hospital admissions to University of Granada Hospital for cryptorchidism from districts near the Mediterranean coast, where there is intensive use of pesticides. A subsequent study reported endosulfan isomers and/or metabolites in adipose tissue of 40% of children who were admitted to the same hospital for a variety of reasons (Olea et al. 1999), indicating that significant endosulfan exposures occurred in the region. In the present study, there is a definite history of endosulfan exposure that is likely to have occurred during the prenatal period.

**Pathways of endosulfan exposure in the study population.** In our study, we estimated endosulfan residues in biologic and environmental samples. The practice of aerial spraying of endosulfan was discontinued in December 2000. Serum endosulfan residue levels were significantly higher in the study population than in the control group even 10 months after the last aerial spray (October 2001). Moreover, endosulfan residues were detected in water ( $\geq 0.03$  ppb) and pond sediments ( $\geq 0.3$  ppb) only in the study area 1.5 years (June 2002) after the last aerial spray. This signifies that low-level endosulfan exposures continued to occur probably by translocation from the hilltops to the valley in the study area long after the aerial spray. This is supported by the report of Regional Remote Sensing Service Center (RRSSC), Bangalore, India (Nageswara Rao PP. Personal communication). On the basis of analysis of satellite pictures of the study area, the RRSSC reported, "The watershed characteristics are favorable for any aerially sprayed toxicant to reach the soil-water-plant continuum in a very short span of time and get accumulated" (Nageswara Rao PP. Unpublished report). Endosulfan has a half life of 60-800 days in soil (ATSDR 2000). Frank et al. (1982) have also reported that because of its persistence in soil, endosulfan residues were detected in water samples throughout the year (outside the spray season) with storm runoff. The results of several laboratory and greenhouse studies indicate that  $\alpha$ - and  $\beta$ -endosulfan are strongly adsorbed to soil (Bowman et al. 1965; El Beit et al. 1981a, 1981b). The study area has an annual rainfall of 140 inches. Twelve first-order streams originate from the cashew plantations. It is likely that endosulfan sticking on the soil is carried by runoff water during most of the year.

We have measured endosulfan levels only once in serum samples. These individual endosulfan measures may or may not accurately reflect the chronic levels and/or levels during critical developmental phases. However, the effect of this would be to decrease power of the study (via an increase in random misclassification of exposure) and thus bias toward the null. We believe that even single estimations of serum endosulfan levels validate that children exposed to endosulfan via aerial spraying do, on average, have higher exposures than children in the control group.

Finally, it is important to discuss and resolve the following weaknesses of the study. First is nonparticipation in SMR study: 57% of the exposed and 33% of the control participants did not agree to undergo SMR examination. However, growth-related end points (height, weight, and skin-fold thickness) were comparable among the nonparticipating groups. It is still difficult to envisage the overall effect of nonparticipation. Second is one-time collection of blood samples for hormone analysis: Sex hormone secretion is pulsatile in nature, with a lot of diurnal variation. To minimize this effect, we collected blood samples in all cases between 1000 and 1200 hr. Because the variation in the blood hormone levels is random, the overall effect of natural variation is likely to be minimized. However, the increased random variability of hormone would have decreased the power of the study via an increase in random misclassification of testosterone levels, which in turn would bias toward the null.



Our study results suggest that endosulfan exposure may delay sexual maturity and interfere with hormone synthesis in male children. To validate these findings, a study with larger sample size is essential. We do not know the significance of these findings. Long-term follow-up of the children is essential to understand the implications.

### References

- [Anonymous.] 2001. Data scan, capacity and technical production of pesticides in India. *Pestic Inf* 17:72-73.
- ATSDR. 2000. Toxicological Profile for Endosulfan. Atlanta, GA:Agency for Toxic Substances and Disease Registry.
- Bowman MC, Schechter MS, Carter RL. 1965. Behavior of chlorinated insecticides in a broad spectrum of soil types. *J Agric Food Chem* 13:360-365.
- Chitra KC, Latchoumycandane C, Mathur PP. 1999. Chronic effect of endosulfan on the testicular functions of rat. *Asian J Androl* 1(4):203-206.
- Dale WE, Curley A, Cueto C Jr. 1966. Hexane extractable chlorinated insecticides in human blood. *Life Sci* 5:47-54.
- Dalsenter PR, Dallegrave E, Mello JRB, Langeloh A, Oliveria RT, Faqi AS. 1999. Reproductive effects of endosulfan on male offspring of rats exposed during pregnancy and lactation. *Hum Exp Toxicol* 18:583-589.
- Dalsenter PR, de Aranjó SL, Silva de Assis HC, Andrade AJM, Dallegrave E. 2003. Pre- and postnatal exposure to endosulfan in Wistar rats. *Hum Exp Toxicol* 22:171-175.
- El Beit IOD, Wheelock JV, Cotton DE. 1981a. Factors affecting soil residues of dieldrin, endosulfan, gamma-HCH, dimethoate, and pyrolan. *Ecotoxicol Environ Saf* 5:135-160.
- . 1981b. Pesticide-microbial interaction in the soil. *Int J Environ Stud* 16:171-179.
- Frank R, Braun HE, Van Hove Holdrinet M, Sirons GJ, Ripley BD. 1982. Agriculture and water quality in the Canadian Great Lakes Basin. V. Pesticide use in 11 agricultural watersheds and presence in stream water, 1975-1977. *J Environ Qual* 11:497-505.
- García-Rodríguez J, García-Martín M, Noguera-Ocaña M, Luna-del-Castillo JD, García MF, Olea N, et al. 1996. Exposure to pesticides and cryptorchidism: geographical evidence of a possible association. *Environ Health Perspect* 104:1090-1095.
- Hack R, Ebert E, Leist KH. 1995. Chronic toxicity and carcinogenicity studies with the insecticide endosulfan in rats and mice. *Food Chem Toxicol* 33:941-950.
- HazDat. 2000. Endosulfan. Atlanta, GA:Agency for Toxic Substances and Disease Registry.
- Khan PK, Sinha SP. 1996. Ameliorating effect of vitamin C on murine sperm toxicity induced by three pesticides (endosulfan, phosphamidon and mancozeb). *Mutagenesis* 11:33-36.
- Marshall WA, Tanner JM. 1969. Variations in pattern of pubertal changes in girls. *Arch Dis Child* 44:291-303.
- Murray TJ, Lea RG, Abramovich D R, Haites NE, Fowler PA. 2001. Endocrine disrupting chemicals: effects on human male reproductive health. *Early Pregnancy* 5:80-112.
- NCI. 1978. 78-Week Dietary Study in Osborne-Mendel Rats and B6C3F1 Mice. NCI Study No. NCI-CG-TR62, Technical Report Series No. 62. Bethesda, MD:National Cancer Institute.
- Olea N, Olea-Serrano F, Lardelli-Claret P, Rivas A, Barba-Navarro A. 1999. Inadvertent exposure to xenoestrogens in children. *Toxicol Ind Health* 15:151-158.
- Sharpe RM. 2001. Hormones and testis development and the possible adverse effects of environmental chemicals. *Toxicol Lett* 120:221-232.
- Singh SK, Pandey RS. 1989. Gonadal toxicity of short term chronic endosulfan exposure to male rats. *Indian J Exp Biol* 27:341-346.
- . 1990. Effect of sub-chronic endosulfan exposures on plasma gonadotrophins, testosterone, testicular testosterone and enzymes of androgen biosynthesis in rat. *Indian J Exp Biol* 28:953-956.
- Sinha N, Adhikari N, Saxena DK. 2001. Effect of endosulfan during fetal gonadal differentiation on spermatogenesis in rats. *Environ Toxicol Pharmacol* 10:29-32.
- Sinha N, Narayan R, Saxena DK. 1997. Effect of endosulfan on the testis of growing rats. *Bull Environ Contam Toxicol* 58:79-86.
- Sinha N, Narayan R, Shanker R, Saxena DK. 1995. Endosulfan-induced biochemical changes in the testis of rats. *Vet Hum Toxicol* 37:547-549.
- U.S. EPA. 1980. Manual of Analytical Methods for the Analysis of Pesticides in Human and Environmental Samples. No. 600/8-80-Section 5A, (3), (a). 038.

Washington, DC:U.S. Environmental Protection Agency.

———. 2002. Re-registration Eligibility Decision for Endosulfan Case No. 0014. Washington, DC:U.S. Environmental Protection Agency.

WHO. 2002. The WHO Recommended Classification of Pesticides by Hazard and Guidelines to Classification 2000-2002. Geneva:World Health Organization, International Programme on Chemical Safety/Inter-Organisation Programme for Sound Management of Chemicals.

---

Last Updated: November 17, 2003

## Science News Online

Week of Dec. 13, 2003; Vol. 164, No. 24

**Slowing Puberty? Pesticide may hinder development in boys**

Kate Ramsayer

Chronic exposure to a widely used pesticide may delay sexual maturity in boys, according to a new study in India.

Endosulfan is an organochlorine used around the world to protect squash, melons, strawberries, and other produce. A 2001 report by the U.S. Environmental Protection Agency estimated that 1.4 million pounds of the pesticide are used annually on U.S. crops. It's also the third most commonly used pesticide in India, where concerns about endosulfan have surfaced.

For example, for irrigation water and fish, the southern Indian town of Padre relies on streams that run through a cashew plantation where the trees had been sprayed with endosulfan a few times a year for 2 decades until December 2000. Residents then demanded that the spraying stop because they suspected it was increasing the local rates of several cancers and neurological diseases.

"Because of the particular topography, people were exposed to high concentrations of endosulfan," says Habibullah Saiyed of the National Institute of Occupational Health in Ahmedabad, India. Even though endosulfan doesn't last long in the body, it can linger in the food chain by contaminating soil and accumulating in fish.

Saiyed and his colleagues collected blood from Padre residents and village-water samples 10 months after the last spray. "They were still showing endosulfan," Saiyed says.

Since previous experiments in laboratory rats had linked high endosulfan exposure during development to defects in sperm quality, the researchers investigated possible effects of the compound on human male-reproductive development. The team examined 117 Padre schoolboys between 10 and 19 years old and a comparable group of boys from a town 20 kilometers away, where there had been no endosulfan spraying.

Blood tests indicated the pesticide's presence in more than three-quarters of the boys in the Padre group, whereas less than a third of the control group's blood samples showed endosulfan. The results demonstrate the overall environmental prevalence of this pesticide, Saiyed says.

Reproductive development of each boy was evaluated using a sexual-maturity rating based on the size of the penis and testes and the development of pubic hair. After taking age into account, the researchers found that boys in Padre scored significantly lower on all three measurements than boys from the other village did. The scientists report their findings in the December *Environmental Health Perspectives*.

They also found that the blood concentration of testosterone was lower in the Padre boys, even though they had higher concentrations of luteinizing hormone than the other boys did. Luteinizing hormone normally pumps up production of testosterone, so the results imply that something is impeding testosterone synthesis and therefore the progression of puberty, the researchers say.

"The findings really suggest that we should use [endosulfan] cautiously or not use it at all," says Saiyed, adding that this is one of the first studies to link an environmental pollutant to delayed puberty in boys.

Richard Sharpe of the Medical Research Council's Centre for Reproductive Biology in Edinburgh says the study ought to raise concerns, but because it looked at only a small number of children, the results may simply represent natural variations in the onset of puberty.

\*\*\*\*\*

If you have a comment on this article that you would like considered for publication in *Science News*, send it to [editors@sciencenews.org](mailto:editors@sciencenews.org). Please include your name and location.

**References:**

Saiyed, H., *et al.* 2003. Effect of endosulfan on male reproductive development. *Environmental Health Perspectives* 111(December):1958-1962. Abstract available at <http://ehpnet1.niehs.nih.gov/docs/2003/6271/abstract.html>.

Information from the U.S. Environmental Protection Agency about endosulfan can be found at [http://www.epa.gov/oppsrrd1/REDS/factsheets/endosulfan\\_fs.htm](http://www.epa.gov/oppsrrd1/REDS/factsheets/endosulfan_fs.htm).

#### **Further Readings:**

Harder, B. 2003. Count down: Chemicals linked to inferior sperm. *Science News* 163(May 31):339. Available at <http://www.sciencenews.org/20030531/fob1.asp>.

Raloff, J. 1998. Common pesticide clobbers amphibians. *Science News* 154(Sept. 5):150. Available at [http://www.sciencenews.org/sn\\_arc98/9\\_5\\_98/fob5.htm](http://www.sciencenews.org/sn_arc98/9_5_98/fob5.htm).

\_\_\_\_\_. 1994. That feminine touch. *Science News* 145(Jan. 22):56. Available at [http://www.sciencenews.org/sn\\_edpik/ls\\_8.htm](http://www.sciencenews.org/sn_edpik/ls_8.htm).

#### **Sources:**

L. Earl Gray  
U. S. Environmental Protection Agency  
NHEERL, Mail Drop 72  
Research Triangle Park, NC 27711

Habibullah Saiyed  
National Institute of Occupational Health  
Meghani Nagar, Ahmedabad - 380 016  
India

Richard Sharpe  
MRC Human Reproductive Sciences Unit  
University of Edinburgh  
Chancellor's Building  
49 Little France Crescent  
Edinburgh EH16 4SB  
United Kingdom

<http://www.sciencenews.org/articles/20031213/fob4.asp>  
From *Science News*, Vol. 164, No. 24, Dec. 13, 2003, p. 372.  
Copyright (c) 2003 Science Service. All rights reserved.

# Endosulfan



January 2001

## What is endosulfan?

Endosulfan is a broad spectrum organochlorine insecticide/ acaricide which is registered in Australia for the control of a large variety of insects and mites in crops. It is not registered for use in animal production.

## Can endosulfan effect my health?

Endosulfan is highly toxic if sufficient exposure occurs by ingestion, skin contact, or inhalation. Lethal poisoning in humans has been reported after accidental or suicidal ingestion of endosulfan. The lowest reported dose that has resulted in death in humans is 35 mg/kg of bodyweight (equivalent to 2.4 grams for a 70 kg adult). Signs of poisoning reported in humans suggest the nervous system is mostly affected. Nausea, vomiting, dizziness, confusion, weakness, hyper-excitability, shortness of breath, decreased respiration, and fine tremors are suggestive of poisoning.

Endosulfan is absorbed through the lungs, skin and gastrointestinal tract. Following absorption, endosulfan (and its metabolites) does not persist for long periods in the tissues or organs and passes relatively quickly from the body.

Endosulfan is not considered to accumulate in the body unlike other organochlorine chemicals (such as DDT).

A number of cases of endosulfan poisoning have been reported in people spraying cotton and rice fields. The dermal route is assumed to be the primary route of exposure. About half the cases presented with convulsions and all cases recovered.

Endosulfan is not considered to be carcinogenic or to adversely affect reproduction or foetal development. There is not any clinical evidence that endosulfan disrupts the hormonal system within the body. Endosulfan is not considered to be an eye irritant or a skin sensitiser. However, it may cause skin irritation.

## How can I be exposed to endosulfan?

In Australia, endosulfan is not registered for use in or around the home. Consequently, the most likely exposure to endosulfan for the public is via residues in food and via accidental exposure (including spray drift).

It is important to recognise that general community (non-occupational) exposure to endosulfan is typically very low and within generally accepted levels. Information from the Australian Market Basket Survey 1996 and studies looking at typical air levels of endosulfan during aerial spraying of cotton crops support this conclusion.



### Occupational Exposure

In occupational settings, exposure to endosulfan is mainly via the dermal and inhalation routes. Exposure levels in occupational settings are generally much higher than those to which members of the public would be exposed. Consequently, to reduce the risk to worker safety that such exposure levels may present, workers need to be aware of regulations relating to the handling of endosulfan. The use of personal safety equipment and/or engineering controls will be required under most circumstances.

### Regulation

Endosulfan is listed in Schedule 7 of the "Standard for the Uniform Scheduling of Drugs and Poisons" on the basis of its acute toxicity.

In Australia, the current Acceptable Daily Intake (ADI) is 0.006 mg/kg/day, based on a no observed effect level (NOEL) of 0.6 mg/kg/day and application of a 100-fold safety factor.


Endosulfan is regarded as a hazardous substance and is covered by regulations to control workplace hazardous substances. Advice from Workplace Health and Safety of the Queensland Government's Department of Employment, Training and Industrial Relations should be sought with respect to issues relating to worker safety.

### Selected References

Toxicological Profile for Endosulfan. Agency for Toxic Substances and Disease Registry, US Department of Health and Human Services, 1998.

The NRA Review Of Endosulfan (August 1998). Volume 1.

For more information, contact your local public health network



Click on the map or the links below to go to contact info

**Central Public Health Unit Network** servicing Brisbane Northside, Longreach, Redcliffe, Rockhampton, Sunshine Coast, Wide Bay.  
For contact details go to <http://www.health.qld.gov.au/phs/cphun/>

**Southern Public Health Unit Network** servicing Brisbane Southside, Darling Downs, Roma, South Coast, South West Queensland, West Moreton  
For contact details go to <http://www.health.qld.gov.au/phs/sphun/>

**Tropical Public Health Unit Network** servicing Cairns, Mackay, Mount Isa and Gulf, Townsville  
For contact details go to <http://www.health.qld.gov.au/phs/tphun/>

For medical advice, contact your doctor, hospital or health clinic.

Published by:  
Environmental Health Unit  
10<sup>th</sup> floor, Queensland Health Building  
147-163 Charlotte Street,  
BRISBANE 4000  
GPO Box 48 BRISBANE 4001  
Ph: +61 7 3234 0938  
Fax: +61 7 3234 1480  
<http://www.health.qld.gov.au/phs/ehu/>

This fact sheet may be reproduced in full, providing the source is acknowledged, but may not be reproduced in part without prior permission of Public Health Services, Queensland Health

